

PROGRAMME DE TRAITEMENT DU PÊCHER-NECTARINE

برنامج علاج أشجار الخوخ و الشهدية



Stades phénologiques du pêcher-nectarine

| A | B | C | D | E | F | J | H | I | G | K | L | M |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | | | | | | | | |

| | | | | | | | | | | | |
|--|------------------|--|--|--|--|--|--|--|--|--|--|
| | Acarie | Apollo 50 SC (40 cc/ha) - Marjoricide (200 cc/ha) | | | | | | | | | |
| | Puceron | Limocide (400 cc/ha) | | | | | | | | | |
| | Thrips | | | | | | | | | | Mavrik 2F (60 cc/ha) Deltaplan 25 EC (30 cc/ha) |
| | Escargot | Cargo 3gb (8 kg/ha) | | | | | | | | | |
| | Cloque du pêcher | Champion (500 gr/ha) - Merjan (500 gr/ha) - Merjan 80 WG (190 gr/ha) | | | | | | | | | |
| | Moniliose | Mancothane 80 (200 gr/ha) | | | | | | | | | |
| | Oïdium | Nimrod 25 EC (60 cc/ha) - Tiosol 800 (160 cc/ha) - Mistral 80 WP (500 gr/ha) | | | | | | | | | |
| | tavelure | Mancothane 80 (200 gr/ha) | | | | | | | | | |
| | Mauvaises herbes | Ovni XL (2 - 4 l/ha) | | | | | | | | | |
| | Nutrition | Ferqueztene (30 à 100 g/arbre) - Maxifer (30 à 100 g/arbre) - Tonofol (400- 500 cc/ha) - Zm Corrector (200-300 g/ha) Bormax foliarel (1 l/ha) - Perfectose (3-6 l/ha) - Blackjak (2-6 l/ha) - Oxicalc (1 l/ha) - Penalty (200-300 cc/ha) - Curane (100 g/ha) Liqui'k Express (300 g/ha) - Top racine (2-3 l/ha) - Stimul (50-60 g/ha) - Corail Ec (3 l/ha) | | | | | | | | | |